


FACTS ARE
STUBBORN THINGS


Rene le Sage

THE COWARDS NEVER
STARTED, AND THE WEAK
DIED ON THE WAY.

Jdit Carson









$$\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

$$Loc_2(R^2) = 3$$

(x_1, y_1)

(x_1, y_1)





$LOC_1(\mathbb{Z} \square \mathbb{Z}) = ?$

branch paths is homeomorphic to $K_{l,n}$. Note that distinct leaves of T must be disjoint. Figure 1 exhibits four topologically distinct trees which have five leaves with stem points labelled v_1, v_2, v_3, v_4 and v_5 . In each of them, v_2 has $\deg(v_2) - 1 = 3$ branch paths. Every tree which is not a path has at least one leaf, and, more generally, the following lemma is easy to prove.

Lemma 1. In a tree T which is not a path, there is one vertex v for which the number of branch paths at v is $\deg(v)$, or there are at least two vertices, v_1 and v_2 , for which the number of branch paths at v_1 and v_2 is $\deg(v_1) - 1$ and $\deg(v_2) - 1$, respectively.






Figure 1. Four trees with the same leaf structure.

- 550 -



$$RED: LOC(G) \leq LOC(G) [1 + 2 \cdot 5^{LOC(G)-1}]$$



$$RED: \gamma(G)/\gamma(G) \rightarrow \infty$$

Fault-Tolerant Metric Dimension of Graphs

83




Figure 1. Tree T' with $\beta(T') = 10$ and $\beta'(T') = 14$.

Theorem 2.2 ([6]). Let T be a tree with set L of endpoints with $|L| \geq 3$. Let L_1, \dots, L_k be the components of the subtree induced by the set of all branch paths, and let e_i be the number of branch paths in T that are in L_i . Then $\beta(T) = |L| - k$, and S is a metric basis if and only if it consists of exactly one vertex from each of exactly $e_i - 1$ of the branch paths of L_i , for each L_i , $1 \leq i \leq k$.

Assume that S is a fault-tolerant resolving set for tree T . For each v in S , the set $S \setminus \{v\}$ is a resolving set. By Theorem 2.1, for any $v \in S \cap L_i$, the set $S \setminus \{v\}$ must contain a vertex from each of $e_i - 1$ of the branch paths of L_i . Thus, $|S \cap L_i| \geq e_i$ when $e_i \geq 2$. If L is the set of endpoints and E_1 is the set of endpoints to branch paths where $e_i = 1$, then, by Theorem 2.1, $L \setminus E_1$ is a fault-tolerant resolving set. This implies the next theorem.

Theorem 2.3. Let T be a tree with set L of endpoints with $|L| \geq 3$. Let L_1, \dots, L_k be the components of the subtree induced by the set of all branch paths, and let e_i be the number of branch paths in T that are in L_i . Let E_1 be the set of endpoints corresponding to branch paths where $e_i = 1$. Then $\beta'(T) = |L \setminus E_1|$ and $L \setminus E_1$ is a fault-tolerant metric basis.

Remark 2.4. If S is a fault-tolerant metric basis for T and s_i is the stem vertex of L_i , then $e_i \geq 3$ implies that $S \cap L_i$ will consist of exactly one vertex from each path of $L_i \setminus \{s_i\}$, and, if $e_i = 2$, then $S \cap L_i$ can be any two vertices of $L_i \setminus \{s_i\}$.

In particular, for the tree T' of Figure 1, $\beta'(T') = 14$ and $(1, 2, 3, \dots, 10, y, v, r, s)$

LOCATING-DOM. $S \subseteq V(G)$


$\forall u, v \in S, N(u) \cap S \neq N(v) \cap S \neq \emptyset$

$$\lceil \frac{2m}{3} \rceil = LD(P_m) = LD(G_m)$$

IHM. $LD(T_m) > m/3$

LD-SET S_1


$|S| \leq \frac{m}{3}$? $S \geq 2$



$$m \leq 3 LD(T_m) - 1$$


$$m \leq LD(G_m) + 2^{LD(G_m)} - 1$$

(RALL, PS [LD-3])



$$m \leq 7 LD(P_m) - 10$$

$$m \leq 7 LD - 10 + 10 \cdot ?$$



$$\beta = |V(G_j) - V(K_x)|$$

$$= 2^x - 1$$

$$j = 2^{x-1}$$

$$i(K_m) = 1 \leq m-1 = LD(K_m)$$

(CHELLALI, MAMOUNI, PS [LD-7])

$$i(T) \leq LD(T)$$

OTHER CLASSES?

S DOM. SHARE $\gamma(n; s) = \sum \frac{1}{1 + n^s}$

$$1 + \frac{1}{2} + \frac{1}{3} = \frac{11}{6}$$

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{6} = \frac{14}{6}$$

$w \in N[a]$

$$\begin{aligned} S \text{ LD } & 1+1+\frac{1}{2}(n-1) \quad LD(G_n) \geq \frac{2}{n+3} n \\ \Rightarrow sl(n) & \leq \frac{n+3}{2} \quad LD\%_0(G^\infty) \geq \frac{2}{n+3} \\ \frac{2}{7} & \leq LD\%_0(\mathbb{Z} \square \mathbb{Z}) = 3/10 \end{aligned}$$

P.J. Slater / Discrete Mathematics 249 (2002) 179–189






Fig. 2. An optimal locating-dominating set for $Z \times Z$.

and all of the values of $1/6$ shifted to y from vertices x (including $x=v$) with $\gamma(x, S) = 7/2$ will be a total of at most $10/3$.

$\exists \text{SAT} \quad \mathcal{U} = \{u_1, \dots, u_N\}$

$\mathcal{C} = \{c_1, c_2, \dots, c_M\} \quad |c_i| = 3$




$LD(G) \leq 5N + M ?$

FTLD

RED: ≤ 1 FAILURE
TO TRANSMIT

FT: ≤ 1 (UNKNOWN WHICH FAILURE TO
DETECT, TRANSMIT ϕ
(ANY $f(v)$ OR $N(v)$ IS TRUE))



THM. $\text{RED:LD}(G) \leq \text{FT:LD}(G)$

THM. \exists a FTLD(G)-set S s.t.
 $N(u) \neq N(v)$

THM. $\text{FTLD}(T_n) > (4/5)n$


THM. $v \in \text{FTLD-set } S$
 $\Rightarrow |\{w \in N(v) : |N(w) \cap S| = 1\}|$

$$\text{Cor. } \delta(v; S) \leq \frac{1}{2} + \frac{1}{2} + \frac{1}{3} / (\deg v - 1) \stackrel{!}{=} \frac{1}{3}$$

$$\text{Cor. } G \text{ r-reg. } \quad \text{FTLD}(G_n) \geq \frac{3}{2+\lambda} \cdot n$$

$$\text{FTLD}_d^*(G^\omega) \geq 3/(2+\lambda)$$

$$\text{Cor. } \text{FTLD}_d^*(ZD2) \geq \frac{3}{2} = 1/2.$$

Fig. 3. $\text{FTLD\%}(Z \times Z) \leq 0.60$.

Theorem 10. $12/23 \leq \text{FTLD\%}(Z \times Z) \leq 3/5$.

Proof. The pattern illustrated in Fig. 3 shows a tiling of $Z \times Z$ using nine vertices in S in each 5-by-3 block, and darkened vertices form an FTLD-set for $Z \times Z$, so $\text{FTLD\%}(Z \times Z) \leq 9/15$.

Let S be an FTLD-set for $Z \times Z$ and $v \in S$. As noted, $\gamma(v; S) \leq 1/2 + 1/2 + 1/3 + 1/3 + 1/3 = 2$. It will be shown that $\gamma(v; S) = 2$ is not possible for any $v \in S$. Assume $v = (i, j)$ and $\gamma(v; S) = 2$. We have $|N[v] \cap S| = 2$ and can assume $(i-1, j) \in S$. Exactly one element $v^* \in N(v)$ contributes $1/2$ to $\gamma(v; S)$, and each of the other three contributes $1/3$.

Assume $v^* = (i-1, j)$. Then $S \cap \{(i-2, j), (i-1, j-1), (i-1, j+1), (i, j-1), (i, j+1), (i+1, j)\} = \emptyset$. Each of $(i, j-1)$, $(i, j+1)$ and $(i+1, j)$ have exactly three neighbors in S , so $\{(i, j-2), (i, j+2), (i+1, j-1), (i+1, j+1)\} \subseteq S$ and $(i+2, j) \notin S$. But now an intruder at $(i+1, j)$ and $(i+1, j-1)$ faulty and an intruder at $(i, j+1)$ and $(i, j+2)$ faulty would both result in (i, j) and $(i+1, j+1)$ reporting a 1 and everything else reporting 0, a contradiction.

Assume $v^* = (i+1, j)$, so $|N[v^*] \cap S| = 2$ and $|N[(i, j-1)] \cap S| = |N[(i, j+1)] \cap S| = |N[(i-1, j)] \cap S| = 3$. If $(i+2, j) \in S$, then $(i+1, j-1) \notin S$, $(i+1, j+1) \notin S$, $(i, j-1)$

LIAR'S LIARS' DOMINATION 8

INTRUDER VERTEX $\alpha \in V(G)$

$$\gamma_{LR}(G) \equiv LR(G) \equiv LR:\gamma(G)$$

$$\equiv \gamma_{LR(1,1)}(G)$$

$LR:\gamma$

$$\gamma_{LR(i,j)}(G)$$

$LR:\gamma$ -SET $L \subseteq V(G)$:
FOR $n \in L$, $\alpha \notin N[n] \Rightarrow n \mapsto \emptyset$

ONE $w \in N[\alpha] \cap L$ CAN HAVE
 $w \mapsto \emptyset$ OR $w \mapsto \gamma \in N[w]$
ELSE $n \in N[\alpha] \cap L \rightarrow n \mapsto \alpha$

- $\gamma_{X(2)}(G) \leq \gamma_{LR}(G) \leq \gamma_{X(3)}(G)$

$u \mapsto s$ and $v \mapsto s \Rightarrow u = v$
 $u \mapsto s, v \mapsto t \neq s, w \in L - \{u, v\}$
 $\Rightarrow w$ TRUTHFUL

- THM $L \subseteq V(G)$ is LR-DOM iff
 - L is $\gamma_{X(2)}$ -DOM
 - $\forall u \neq v, |L \cap (N[u] \cup N[v])| \geq 3$



- COR EACH COMPONENT OF $\langle L \rangle$ HAS $|el| \geq 3$.

THM. $\Delta(G) = \lambda \Rightarrow LR(G) \geq \frac{6}{3\lambda+2} \cdot n$

R. (COMPNT SIZES ≥ 3 and $\delta_{X(2)}$ SUFFICES)

ASSUME $LR(G)$ -set L , $|L| = k$


$$\sum_{v \in L} \deg v \leq k\lambda$$

$e \in L$ contributes 2 to \sum


so $|E(L, V(G) - L)| \leq k\lambda - 2(\frac{2}{3}\lambda)$

 $\Rightarrow n \leq k + \frac{1}{2}[k(\lambda - 4/3)]$
 $= k(\frac{3\lambda+2}{6}) \Rightarrow k \geq \frac{6}{3\lambda+2} \cdot n$

THM. $\frac{3}{4}(n+1) \leq LR(T_m) \leq n$



v : ENDPOINT
 v : $T-v$ HAS
SOME $|C| \leq 2$



$$\gamma_{LR}(G) \equiv \gamma_{X(2,3)}(G)$$

SET-SIZED DOMINATION


$$\gamma_{X(c_1, c_2, c_3, \dots, c_k, \dots)}$$

LPPDOM : Is $\gamma_{X(2,3)}(G) \leq j$?

DOM : Is $\gamma(G) \leq k$?

DOM \propto LR: $\gamma \equiv \gamma_{X(2,3)}$

$\gamma_{X(c_1, c_2)}(G) \leq j$?
NP-C.



(03) (13) (23) (3)

; ; ;

(02) (12) (2)

(1,3) (2,3) (3,3) ...

(01) (1)

(1,2) (2,2) (3,2) ...

(00)

(1,1) (2,1) (3,1) ...

$\gamma_{X(i,j)}$

LR(i,j)

$LR(i,j) \equiv \gamma_{X(j+1, 2j+1)}$

$LR(1,1) \neq \gamma_{X(c_1, c_2, \dots)}$

$S = \{S_1, S_2, \dots, S_k\}$ IS
DISTINGUISHING FOR G IF

- $\cup S_i = V(G)$
- $\forall u \neq v \exists S_i \ni |S_i \cap \{u, v\}| = 1$

$N[v_i] \neq N[v_j]$ IC $\{N[v_1], N[v_2], \dots, N[v_k]\}$

$N(v_i) \neq N(v_j)$ OLD $\{N(v_1), N(v_2), \dots, N(v_m)\}$

LOC $\{\underline{\{v_1\}}, \underline{N(v_1)}, \underline{N^2(v_1)}, \dots,$
 $\underline{\{v_m\}}, \underline{N(v_m)}, \underline{N^2(v_m)}, \dots\}$

LD $\{\underline{\{v_1\}}, \underline{N(v_1)}, \underline{\{v_2\}}, \underline{N(\{v_2\})}, \dots$
 $\underline{\{v_m\}}, \underline{N(N_m)}\}$

Y $\{\underline{\{v_{1,1}\}}, \dots, \underline{\{v_{1,1+d_1}\}}, \underline{\{v_{2,1}\}},$
 $\dots, \underline{\{v_{m,1}\}}, \dots, \underline{\{v_{m,1+d_m}\}}\}$

S $S(v) = \{i \mid v \in S_i\} \neq \emptyset$
 $|S(u) \Delta S(v)| \geq 1$

S is REDUNDANT DISTING.
FOR G if all $S - \{S_i\}$ are DISTING.
 $\Rightarrow "S_{x(2)}$

$$S \quad S(G) \geq 1$$

$$LD \quad S(G) \geq 1$$

$$LOC \quad S(G) \geq 1$$


$$IC \quad |N[u] \Delta N[v]| \geq 2$$

$$OLD \quad |N(u) \Delta N(v)| \geq 2 \quad OLD = 3$$


$$S: |S(u)| \geq 2 \quad |S(u) \Delta S(v)| \geq 2$$

" $S_{x(2)}$ "

$$IC = 3$$



#



#

\mathcal{A} is FT-DISTINGUISHING if \mathcal{A} CAN DISTING. UNDER THE CONDITION THAT ONE s_i CAN FALSELY REPORT \emptyset .

FT:IC $s \leq v(f)$

- s DOUBLE DOMINATES
- \forall PAIR $u, v \exists x, y \in s$
 $\exists \phi \neq N[x] \cap \{u, v\} = N[y] \cap \{u, v\} \neq \{u, v\}$

\mathcal{A} is FT-DISTING.:

- $\forall u$ WE HAVE $|A(u)| \geq 2$
- $\forall u, v \exists i, j$ WITH
 $\phi \neq s_i \cap \{u, v\} = s_j \cap \{u, v\} \neq \{u, v\}$

EITHER $|A(u) \setminus A(v)| \geq 2$

OR $|A(v) \setminus A(u)| \geq 2$

$\gamma \leq \text{RED} : \gamma = \text{FT} : \gamma \leq \text{LR} : \gamma \leq \text{ER} : \gamma$ $\gamma_{x(2)}$ $\gamma_{x(2,3)}$ $\gamma_{x(3)}$  $\gamma \frac{1}{3}$  $\text{RED} : \gamma \equiv \gamma_{x(2)}$ $\frac{2}{3}$ $\text{FT} : \gamma \equiv \gamma_{x(2)}$ $\frac{2}{3}$  $\text{LR} : \gamma \equiv \gamma_{x(2,3)}$ $\frac{3}{4}$  $\text{ER} : \gamma \quad 1$

IF $|S \cap N[w]| = |\{\alpha, \alpha\}|$

AND $\alpha \mapsto w, \alpha \mapsto \emptyset$

THEN INTRUDER AT w

OR \nexists AN INTRUDER

 $\text{ER} : \gamma \equiv \gamma_{x(3)}$ $\text{ER}^- : \gamma$

$LD \leq RED:LD \leq FT:LD \leq LR:LD \leq ER:LD$

$y_{x(2)} \leq$



$LD \quad 2/5$



$RED:LD \quad 2/3$



$FT:LD \quad 4/5$



$LR:LD \quad 4/5$



$ER:LD \quad 1$

$$LOC \leq RED:LOC \approx FT:LOC \leq LR:LOC \\ = ER:LOC$$



LOC $LOC(P_m) = LOC(P_{\infty}) = 1, LOC(P_\infty) = 2$

$RED:LOC(P_m) = 2$ $RED:LOC(P_\infty) = RED:LOC(P_\infty) = 3$

$FT:LOC \leq RED:LOC$



$LR:LOC = 4$

$LR:LOC(P_m) = 4, m \geq 4$

For $m \geq 4$,


$3 \leq LR:LOC(G_m) = ER:LOC(G_m)$

$IC \leq RED : IC \leq FT : IC \leq LR : IC \leq ER : IC$




$IC = \frac{1}{2}$

$RED : IC = "1"$



$RED : IC(G) \neq 1 \text{ if } \exists w \in$

$W(G) - w \text{ has a 2-component}$



$IC = 2\lambda$

$RED : IC = 3\lambda = n$




$FT : IC$



001100

$u \text{ OR } n$

$FT : IC(T) \neq \text{tree } T$



$LR : IC \equiv FT : IC$

DOMINATION LOC-PON IC OLD LOC ...

REDUNDANCE: FAILURE TO TRANSMIT

RED:S RED:LD RED:IC RED:OLD RED:Loc

FAULT TOLERANCE: UNKNOWN FAILURE
TO DETECT

FT:S FT:LD FT:IC FT:OLD FT:Loc

LIAR'S: MISREPRESENTING A / SET
POSITIVE DETECTION SIZED
POMWATH

LR:S LR:LD LR:IC LR:OLD LR:Loc

ERROR: ANY MISREPRESENTING

ER:S ER:LD ER:IC ER:OLD ER:Loc